




Redx

Discovering Targeted Medicines

Corporate Presentation

February 2022

AIM:REDX



Disclaimer



This presentation has been prepared by Redx Pharma plc (the "Company"). By reviewing this presentation you agree to be bound by the following conditions.

The information being supplied in this presentation is being supplied and communicated to you solely for your information and may not be reproduced or further distributed to any person or published, in whole or in part, for any purpose.

The presentation has not been independently verified and no representation or warranty, express or implied, is made or given by or on behalf of the Company or any of its parent or subsidiary undertakings, or the subsidiary undertakings of any such parent undertakings, or any of such person's respective directors, officers, employees, agents, affiliates or advisers, as to, and no reliance should be placed on, the accuracy, completeness or fairness of the information or opinions contained in this presentation and no responsibility or liability is assumed by any such persons for any such information or opinions or for any errors or omissions. All information presented or contained in this presentation is subject to verification, correction, completion and change without notice. In giving this presentation, none of the Company or any of its parent or subsidiary undertakings, or the subsidiary undertakings of any such parent undertakings, or any of such person's respective directors, officers, employees, agents, affiliates or advisers, undertakes any obligation to amend, correct or update this presentation or to provide the recipient with access to any further information that may arise in connection with it.

Neither this presentation nor anything contained herein shall form the basis of or be relied upon in connection with, or act as an inducement to enter into, any investment activity. This presentation does not purport to contain all of the information that may be required to evaluate any investment in the Company or any of its securities and should not be relied upon to form the basis of, or be relied on in connection with, any contract or commitment or investment decision whatsoever. This presentation is not intended to provide complete disclosure upon which an investment decision could be made. The merit and suitability of an investment in the Company should be independently evaluated and any person considering such an investment in the Company is advised to obtain independent advice as to the legal, tax, accounting, financial, credit and other related advice prior to making an investment.

This presentation is made available for information purposes only and does not, and is not intended to, constitute an offer to sell or an offer, inducement, invitation or commitment to purchase or subscribe for any securities. The distribution of this presentation may, in certain jurisdictions, be restricted by law and persons into whose possession this presentation comes should inform themselves about, and observe, any such restrictions. Neither this presentation, nor any part of it nor the fact of its distribution shall form the basis of or be relied upon in connection with any contract and it does not constitute a recommendation regarding any securities.

To the extent available, the data contained in this presentation has come from official or third party sources. Third party industry publications, studies and surveys generally state that the data contained therein have been obtained from sources believed to be reliable, but that there is no guarantee of the accuracy or completeness of such data. While the Company believes that each of these publications, studies and surveys has been prepared by a reputable source, the Company has not independently verified the data contained therein. In addition, certain of the data contained in this presentation come from the Company's own internal research and estimates based on the knowledge and experience of the Company's management in the market in which the Company operates. While the Company believes that such research and estimates are reasonable and reliable, they, and their underlying methodology and assumptions, have not been verified by any independent source for accuracy or completeness and are subject to change without notice. Accordingly, undue reliance should not be placed on any of the data contained in this presentation.

This presentation has not been approved by an authorised person in accordance with section 21 of the Financial Services and Markets Act 2000. As such, this document is being made available only to and is directed only at: (a) persons outside the United Kingdom, (b) persons who have professional experience in matters relating to investments who fall within the definition of "investment professionals" in Article 19(5) of the Financial Services and Markets Act 2000 (Financial Promotion) Order 2005 (the "FPO"), (c) high net worth bodies corporate, unincorporated associations and partnerships and trustees of high value trusts as described in Article 49(2)(a) to (d) of the FPO or (d) those persons to whom it may otherwise be lawfully communicated (in each case referred to as "Relevant Persons"). The information contained in this document is not intended to be viewed by, or distributed or passed on (directly or indirectly) to, and should not be acted upon by any class of persons other than Relevant Persons. Any investment or investment activity to which this document relates is available only to Relevant Persons and will be engaged only with Relevant Persons.

This presentation includes forward looking statements. The words "expect", "anticipate", "intends", "plan", "estimate", "aim", "forecast", "project" and similar expressions (or their negative) identify certain of these forward looking statements. These forward looking statements are statements regarding the Company's intentions, beliefs or current expectations concerning, among other things, the Company's results of operations, financial condition, liquidity, prospects, growth, strategies and the industry in which the Company operates, and include statements regarding the Company's ongoing and planned clinical trials, regulatory approval process, and demand for the Company's product candidates which are subject to risks, uncertainties, and other factors that could cause actual results to differ materially from those suggested by forward-looking statements. These factors include, but are not limited to, the following: the Company has incurred losses since its inception and anticipates that it may continue to incur losses for the foreseeable future; the Company does not expect to generate any material income until its pipeline of programmes are progressed commercially; the Company will need to raise additional funding in the future, which may not be available on acceptable terms, or at all; and the Company may not be able to obtain exclusivity or intellectual property rights for its product candidates or prevent others from developing similar competitive products. The forward looking statements in this presentation are based on numerous assumptions regarding the Company's present and future business strategies and the environment in which the Company will operate in the future. Forward looking statements involve inherent known and unknown risks, uncertainties and contingencies because they relate to events and depend on circumstances that may or may not occur in the future and may cause the actual results, performance or achievements of the Company to be materially different from those expressed or implied by such forward-looking statements. Many of these risks and uncertainties relate to factors that are beyond the Company's ability to control or estimate precisely, such as future market conditions, currency fluctuations, the behaviour of other market participants, the actions of regulators and other factors such as the Company's ability to continue to obtain financing to meet its liquidity needs, changes in the political, social and regulatory framework in which the Company operates or in economic or technological trends or conditions. Past performance should not be taken as an indication or guarantee of future results, and no representation or warranty, express or implied, is made regarding future performance. The Company expressly disclaims any obligation or undertaking to release any updates or revisions to these forward looking statements to reflect any change in the Company's expectations with regard thereto or any change in events, conditions or circumstances on which any statement is based after the date of this presentation or to update or to keep current any other information contained in this presentation. Accordingly, undue reliance should not be placed on the forward looking statements, which speak only as of the date of this presentation.

Ambitious Leadership Team with Deep Industry Experience



Lisa Anson

Chief Executive Officer

>25 years biotech and pharma, with significant senior level commercial leadership and general manager experience

Sector profile as BIA Board member and prior President of ABPI

MBA, MA Natural Sciences



Dr Richard Armer

Chief Scientific Officer

>25 years biotech and pharma with significant translational science experience bringing products from discovery to the clinic

Experience at Pfizer, Organon, Ardana, Oxagen and Lectus Therapeutics

PhD in Chemistry



Dr Jane Robertson

Chief Medical Officer

>18 years in biotech and pharma, with extensive experience as former CMO at Achilles & Nucana and leading oncology clinical development programs including Lynparza

13 years in clinical practice

MD, MRCP, FRCPath



Peter Collum

Chief Financial Officer

Experienced biopharma finance and strategy executive with >24 years in biopharma as CFO, CBO and healthcare investment banking

MBA, BS in Engineering



Dr James Mead

Chief Operating Officer

>20 years in biotech and pharma with multiple finance leadership roles including CFO positions as well roles covering Investor Relations and corporate development

PhD in molecular biology, Chartered Accountant



Claire Solk

General Counsel

Experienced legal counsel with wide sector experience

Previously with AstraZeneca and Allen & Overy LLP

Admitted to practise in England & Wales and New York





Novel, small molecule, highly targeted medicines for cancer and fibrotic disease, including the emerging area of cancer-associated fibrosis



Clinical-stage biotech with two wholly-owned clinical assets and Discovery Engine fuelling preclinical pipeline



Ambitious, experienced leadership/scientific team with distinct approach and track record for consistently generating differentiated drug candidates



Pipeline financed to deliver multiple value inflection points in 2022 and backed by blue-chip specialist biotech investors

A Clinical Stage Biotech with Three Value Drivers

Selective Porcupine Inhibitor (RXC004)

- Ph1 monotherapy: differential activity in Wnt-ligand dependent tumours
- Ph1 combo ongoing with nivolumab
- Ph2 (PORCUPINE) initiated: genetically selected MSS mCRC
- Ph2 (PORCUPINE2) initiated H1 2022: genetically selected pancreatic/unselected biliary
- Open IND in the US

Selective ROCK2 Inhibitor (RXC007)

- Targeting a master switch in fibrosis
- Clinically validated target, with first ROCK2 inhibitor approved by FDA in 2021
- Interim Ph1 data: no adverse events (up to 40 mg as a single dose)
- Biologically relevant exposures
- Lead indication IPF – Ph2 start in 2022

Discovery Engine

- Strong track record:
 - Generated four molecules now in clinic incl. pirtobrutinib
 - Partnership deals with Jazz and AstraZeneca
- Partnered programmes: \$26m milestones earned in 2021
- Discoidin domain receptor (DDR) inhibitors for fibrosis
- Targeting 3 wholly-owned INDs by 2025

IND: Investigational New Drug application; IPF: Idiopathic Pulmonary Fibrosis; ROCK: Rho Kinase; Wnt: Wingless/integrated


Robust Pipeline with Two Wholly-Owned Clinical Stage Assets Built from In-House R&D Engine



	Target/Product	Research	Preclinical	Phase 1	Phase 2	Phase 3	Upcoming Milestones
Oncology	Porcupine Inhibitor (RXC004)	Genetically selected MSS mCRC mono + nivolumab combo					Combo: Initiate Phase 2 - 2022 Topline data - H1 2023
		Genetically selected pancreatic cancer Unselected biliary cancer					Initiate Phase 2 - H1 2022 Topline data - H1 2023
Fibrosis	ROCK2 Selective Inhibitor (RXC007)	Idiopathic pulmonary fibrosis (IPF)					Phase 1 topline results - H1 2022 Initiate Phase 2a - 2022
	GI-targeted ROCK Inhibitor	Fibrosis in Crohn's					Preclinical development candidate - H1 2022
	DDR Inhibitor (Discoidin Domain Receptor)	Fibrosis					In lead optimisation
Partnered	MAPK Pathway (2 targets)	Oncology					Progress Jazz collaboration
	Porcupine Inhibitor (RXC006/AZD5055)	Idiopathic pulmonary fibrosis (IPF)					Licensed to AstraZeneca
	Pan-RAF Inhibitor (JZP815)	Oncology					Sold to Jazz Pharmaceuticals

DDR: Discoidin Domain Receptor; RAF: Rapidly accelerated fibrosarcoma; GI: Gastrointestinal; MAPK: Mitogen-activated protein kinase; MSS mCRC: Microsatellite-Stable Metastatic Colorectal Cancer; ROCK: Rho associated protein kinase

RXC004: Clinically Validated Porcupine Inhibitor in Phase 2



RXC004 is a potent, orally active Porcupine Inhibitor

- Differentiated vs. competitors
- Encouraging Phase 1 monotherapy data reported at ESMO 2021, efficacy signal consistent with activity in Wnt-ligand selected tumours

Porcupine is a key enzyme in the Wnt pathway

- Wnt pathway is well established as a critical driver of cancer
- Inhibition of Porcupine blocks the release of all Wnt ligands from cells, **preventing both tumour growth and tumour immune evasion**
- Clear patient selection - targeting **Wnt-ligand dependent** tumours

Phase 2 programme data expected H1 2023

- **Monotherapy and combination** in genetically-selected MSS metastatic colorectal cancer
- Genetically selected pancreatic cancer
- Unselected biliary cancer

ESMO: European Society for Medical Oncology; RNF43: Ring Finger Protein 43; RSPO: R-Spondin; Wnt: Wingless/integrated

Phase 1: Monotherapy Complete; Data from Dose Escalation for Anti-PD1 Combination Expected H1 2022



Phase 1 Objective: Assess safety and tolerability of RXC004 in all-comer cohorts of advanced cancer patients

Dose escalation - single ascending dose / multiple ascending dose, 3+3 design


Module 1: Monotherapy, 0.5-3.0 mg, N=24

Complete; presented at ESMO 2021

- Drug well tolerated in patients up to 2 mg as monotherapy
- Most common AEs: fatigue, decreased appetite, nausea, dysgeusia, vomiting
 - Dysgeusia appears dose-related; did not lead to discontinuations
- 2 mg selected as Phase 2 monotherapy dose

Module 2: Nivolumab Combination, 1.0-2.0 mg

Ongoing



- 1 mg dose was well tolerated in combination with nivolumab 480 mg every four weeks


Module 3: Monotherapy Dose Scheduling, 2 mg, N=12

Initiating 2022

Clinical Efficacy Data Supports Development in Wnt-Ligand Dependent Tumours



Clinical Activity by Wnt-Ligand Dependence†



18 patients had RECIST - evaluable disease

5/7 patients with Wnt-ligand dependent tumours had durable RECIST Stable Disease (SD)

- In addition, 2/7 patients with Wnt-ligand dependent tumours had SD in target lesions but disease progression overall
- 0/11 patients with unknown or Wnt-ligand independent tumours had SD

Median treatment duration higher in patients with Wnt-ligand dependent tumours

- 13.1 weeks in Wnt-ligand dependent tumours
- 6.6 weeks in unknown or Wnt-ligand independent tumours


APC: Adenomatous Polyposis Coli Gene; CRC: Colorectal Cancer; LoF: Loss of Function; PD: Progressive Disease; RECIST: Response Evaluation Criteria in Solid Tumours; RNF43: Ring Finger Protein 43; RSPO: R-Spondin; SD: Stable Disease; Wnt: Wingless/integrated

Data cut-off date 30 July 2021

Data in Genetically-Selected Cancer Models Supports Potential for Enhanced Responses in Selected Patients




Cells with RNF43 mutations or RSPO fusions are sensitive to RXC004 *in vitro*



Inhibition of cell proliferation with RXC004 across 11 genetically-defined cancer cell models of Wnt pathway aberration

Efficacy translates *in vivo* in genetically defined xenograft models




RNF43 mutant human pancreatic cancer line (Capan-2) grown in a mouse xenograft model (10 mice/group)

Preclinical Data Indicates Wnt Pathway Blockade Could Enhance Immune Related Survival - Supporting Combination Development



RXC004 is efficacious in a "cold" tumour model



Improved survival rate observed as monotherapy in B16F10 syngeneic melanoma immune mediated model


Anti-PD1 had no monotherapy effect on this immunologically "cold" model

- Immune-checkpoint inhibitors (ICI) ineffective in MSS mCRC (~95% MSS mCRC⁽¹⁾)
- Wnt driven tumour immune evasion: Wnt activation leads to ICI resistance across 28 cancer types^(2,3)
- RXC004 potential to initiate immune responses in "cold" tumours, where anti-PD1 ineffective, and "hot" tumours to improve ICI responses⁽⁴⁾
- Porcupine inhibitor (WNT974, Novartis) in combination with ICI in ongoing clinical trial demonstrates acceptable safety and early proof of concept⁽⁵⁾

(1) Gong et al. March 21, 2017 (ASCO JCO); (2) Spranger et al., 2015; (3) Luke et al., 2019; (4) Phillips et al 2019; (5) Janku et al. (2020) AACR, CT034 - Phase I study of WNT974 + spartalizumab in patients with advanced solid tumors;

* Survival data calculated from the point at which mice were sacrificed when they reached 2500mm³ tumour volume
ICI: immune-checkpoint inhibitor, mCRC: metastatic colorectal cancer; MSS: Micro Satellite Stable

Phase 2 Programme in Wnt-ligand Dependent Tumours Expected Redx to Deliver Topline Data H1 2023



⁽¹⁾RNF43 mutation frequency determined from all relevant studies published on cBioPortal for cancer genomics (updated Jan 2018). Only mutations resulting in functional impairment (LoF) were considered. Gao et al. 2013 & Cerami et al. 2012. ⁽²⁾RSPO fusion prevalence in CRC is a combination of studies (Shesagiri, 2012; Shinmura, 2014; Kleeman, 2019) ; ⁽³⁾Loilome et al. 2014, Boulter et al. 2015; ⁽⁴⁾www.cancer.net ⁽⁵⁾Incidence data sourced from GlobalData Epidemiology data (Major Markets: US, EU5, Japan, China) ⁽⁶⁾Gong et al. March 21, 2017 (ASCO JCO) ⁽⁷⁾Witkiewicz 2015.
MSS mCRC: Microsatellite-Stable Metastatic Colorectal Cancer; RNF43: Ring Finger Protein 43; RSPO: R-Spondin

Significant Opportunity in Genetically Selected Wnt-ligand Driven Cancers with High Unmet Need



Approximately 74,000 patients per year in the initially targeted indications

	5 Year Survival (metastatic disease)	Annual incidence of (new) metastatic cases	Prevalence of Wnt-driven tumours		Annual target market (new patients)
MSS mCRC 	15%	178k	8% (3% RNF43 + 5% RSPO)	>	14k
Pancreatic Cancer 	3%	130k	7% (RNF43)	>	9k
Biliary Cancer 	2%	>51k	70% (>70% of all biliary cancer Wnt-driven)	>	>51k

* Source: GlobalData from 7 major markets (US, EU5, Japan)


RXC004 Showed Clinical Efficacy in Wnt-Ligand Dependent Tumours and Manageable Toxicity in Phase 1



Highlights

- Porcupine inhibition blocks all Wnt ligands, preventing tumour growth and immune evasion
- RXC004 is a highly potent, orally active, once daily Porcupine inhibitor
- Phase 1 clinical efficacy data supports development in Wnt-ligand dependent tumours
- RXC004 monotherapy was well tolerated at doses up to 2mg
- Target engagement evident at all doses, with doses $\geq 1.5\text{mg}$ achieving exposures that demonstrated efficacy in all preclinical models tested
- Selected dose for Phase 2 monotherapy studies is 2mg
- Phase 2 monotherapy programme initiated in Q4 2021; combination with nivolumab expected to commence in 2022

RXC007 Is a Selective ROCK2 Inhibitor for Fibrotic Diseases - Phase 1 Topline Data Expected H1 2022



RXC007 is a highly selective oral ROCK2 Inhibitor

- Compelling preclinical efficacy demonstrated across fibrotic disease models
- Phase 1 topline data expected in H1 2022

ROCK2 is a compelling fibrosis target

- ROCK sits at a key junction in cell signalling pathways central to fibrosis
- Systemic inhibition of ROCK1&2 results in hypotension, not seen with ROCK2 selective inhibition
- Validated target with a product approved by FDA for cGvHD


Phase 2a in IPF expected to commence 2022

- Idiopathic Pulmonary Fibrosis (IPF) high unmet need - median survival 3-5 years
- IPF market projected to reach \$3.6 bn by 2029*

* Global Data (US, EU5, Japan)

cGvHD: Chronic graft versus host disease; ROCK: Rho associated protein kinase

RXC007 is a Highly Potent and Selective ROCK2 Inhibitor




- ROCK has been a notoriously difficult target
- Systemic inhibition of ROCK1&2 results in hypotension, not seen with ROCK2 selective inhibition
- Need high selectivity for safety
- RXC007 is highly selective, >100-fold versus ROCK1 in biochemical assays

Strong Safety and Pharmacokinetic Profile in Phase 1 in Healthy Volunteers



PK profile after single dose administration over 24 hours*



- No adverse events observed to date following single oral dose of 2-40 mg
- Pharmacokinetics as predicted from preclinical data
- Essentially linear exposure 2-40 mg
- Achieved biologically relevant exposures at 20mg and above
- At 40 mg, half-life ($t_{1/2}$) ~11 hours, potentially suitable for once-daily dosing
- Single ascending dose (SAD) escalations ongoing
- **Phase 1 results expected in H1 2022**



* Interim data on 14 September 2021

Redx internal data

PK: Pharmacokinetic

Good activity in industry standard IPF therapeutic model (bleomycin)



Reduction of fibrosis and collagen deposition



Representative images of lung morphology

Strong modulation of key inflammatory and fibrotic gene signals



Human post-transplant IPF tissue



Human precision cut lung slices (PCLS) mRNA analysis

Activity in Multiple Organs

Reduction of fibrosis and collagen deposition in CCl₄ (carbon tetrachloride) therapeutic model of liver fibrosis




Representative images of liver morphology

Strong Preclinical Data Supports IPF as Lead Indication



Efficacy in well-validated bleomycin *in vivo* model of lung fibrosis




Values expressed as mean of n = 5-11 + SEM. Uncorrected Fisher's LSD model. No differences at significance level 0.05, *: P < 0.05, **: P < 0.01, ***: P < 0.001, ****: P < 0.0001 compared to Bleomycin Vehicle † Pirfenidone dose of 100 mg/kg BID corresponds to a supra-therapeutic dose in humans

Phase 2 Programme in IPF Expected to Commence in 2022




Phase 2a Dose Ranging Study to inform Phase 2b dose; 3 cohorts of 16 patients each

- Provides early efficacy readouts, safety and tolerability in IPF patients, with or without standard IPF therapy
- Assigned dosing period of 3 months. Patients may continue for longer if no signs of disease progression



Translational Science Sub-Study to evaluate target and disease marker engagement; 2 cohorts of 8 patients each




Significant Market Opportunity in IPF



Normal Lung



IPF Lung



- IPF is a deadly fibrotic disease, median survival 3 – 5 years ⁽¹⁾
- 170,000+ patients ⁽²⁾
- IPF market projected to reach \$3.6 billion by 2029 ⁽²⁾
- Nintedanib and pirfenidone are the only approved treatments for IPF
 - Slow the progression of the disease
 - Side effects that limit use
 - Clear opportunity to improve on standard of care

1. Clinical Estimates from Hyun 2015, Ley 2012, Raghu 2006

2. Patient numbers (diagnosed prevalence) & market size forecast data sourced from Global Data (US, EU5, Japan)

IPF: Idiopathic Pulmonary Fibrosis

RXC007 Clinical Development Planned in IPF with Staged Phase 2 Start Expected 2022




Highlights

- ROCK is a validated compelling target at a key junction in cell signalling pathway central to fibrosis
- RXC007 is a highly potent, selective and orally active ROCK2 inhibitor
- Strong preclinical efficacy data supports clinical development plan in IPF
- To date* no adverse events observed in Phase 1 healthy volunteers and observed PK profile in line with prediction from preclinical data
- Dose escalations ongoing in Phase 1
- Plan to start a staged Phase 2 approach in 2022
- Initial 12-week Phase 2a study for early efficacy readouts, safety and tolerability in IPF patients +/- SoC, in addition to target and disease biomarker engagement
- Phase 2a dose-ranging study will inform Phase 2b dose
- Phase 2b will dose RXC007 over 12 months plus SoC with FVC lung function primary endpoint

*Interim data at 14 September 2021

FVC: Forced Vital Capacity; IPF: Idiopathic Pulmonary Fibrosis; PK: Pharmacokinetic; ROCK: Rho Kinase; SoC: Standard of care

Validated Discovery Engine Fuels Pipeline



ADME: Absorption, distribution, metabolism and excretion; IND: Investigational New Drug application; PoC: Proof of Concept; PK: Pharmacokinetic

GI-targeted ROCK Inhibitor Targeting Drug Candidate in 2022




Inflammatory




Fibrostenotic




Fistulizing



Untreated



2.5% DSS 9 wk



DSS + GI-targeted ROCK
inhibitor 3mg/kg QD

Potent, oral small molecule ROCK 1/2 inhibitor

- ROCK is a key target involved in fibroblast activation
- Selectively active in gut without risking systemic exposure
- *In vivo* efficacy in models and *ex vivo* using tissue from Crohn's patients

Potential first-in class treatment for Crohn's disease-associated fibrosis (fibrostenosis)

- Crohn's disease affects 1.5m¹ people globally, of which 50% will develop strictures or complications leading to fibrostenosis²
- No treatment is currently available except invasive surgery
- No experimental therapies for underlying fibrosis


Preclinical development candidate selection in 2022

GI-targeted ROCK inhibitor reduces collagen in mouse model of Crohn's fibrosis

Increase of collagen staining shown in blue in the DSS treated animals. GI-targeted ROCK inhibitor reduces production of collagen seen as a reduction in blue (trichrome) staining.

(1) GlobalData Crohn's Disease Dynamic Market Forecast to 2026 report; (2) Chan et al, 2018;

Discoidin Domain Receptor (DDR) Inhibitors as a Potential Novel Therapeutic Class for Fibrosis



DDR is a collagen target

- Non-integrin tyrosine kinase collagen receptors
- Collagen binding initiates downstream fibrotic signalling pathways

DDR inhibition is a novel approach

- Novel, druggable therapeutic target for fibrosis
- Recent publication of pre-clinical PoC studies for small molecule inhibitors in models of lung and kidney fibrosis
- Potential for disease modifying efficacy

Redx has a discovery programme

- Potent and selective DDR inhibitors identified
- In lead optimisation phase
- Pre-clinical PoC ongoing

Strong Financial Position with Support from Top Tier Specialist Investors



Strong Financial Position

£29.6 million cash

as at 30 September 2021

Cash runway through 2022*

Productive Collaborations

\$26 million in milestones earned in 2021

~\$1 billion in additional potential milestones



Top Tier Specialist Investors

Redmile Group



SOFINNOVA
PARTNERS



POLAR CAPITAL



AIM (UK) listed Ticker: REDX
Total shares in issue: 275,282,205
Fully diluted: 419,148,197**

* Cash through Q4 2022 calendar year.

Multiple Potential Milestones During 2022



Selective Porcupine Inhibitor (RXC004)

- H1 2022 - Data from Phase 1 combination study with nivolumab (anti-PD1)
- H1 2022 - Initiation of PORCUPINE2 Phase 2 study in genetically selected pancreatic and unselected biliary cancer
- 2022 - Initiation of PORCUPINE combination arm in genetically selected MSS mCRC

Selective ROCK2 Inhibitor (RXC007)

- H1 2022 - Data from Phase 1 healthy volunteer study
- 2022 - Initiation of Phase 2a study in IPF
- H2 2022 - US IND

Discovery Engine


- 3 wholly-owned INDs by 2025
 - Targeting DC for GI-targeted ROCK inhibitor for Crohn's fibrosis
 - Progress discoidin domain receptor (DDR) inhibitor for fibrosis
 - Advance multiple programmes in early discovery
- Partnered programmes progress towards next milestones

GI: Gastrointestinal; IND: Investigational New Drug application; IPF: Idiopathic Pulmonary Fibrosis; LO: Lead optimisation; MSS mCRC: Microsatellite-Stable Metastatic Colorectal Cancer; ROCK: Rho Kinase; Wnt: Wingless/integrated
All dates listed are calendar year


Appendix



Inhibits Tumour Cell Proliferation




Control




+RXC004

Reduces
Ki67 staining in
RSPO-fusion CRC
xenograft model

Differentiates Tumour Cells




Control



+RXC004

Increased size and mucin staining of tumour cells with RSPO-fusion. Cells also less metabolically active


As a Monotherapy



Increased expression following RXC004


Increase in
immune related
gene expression
within tumours post-
RXC004 treatment in
anti-PD1-resistant
mouse melanoma
syngeneic model
(B16F10)

In Combination with anti-PD1



Improved responses observed when RXC004 combined with anti-PD1 in syngeneic CRC immune-mediated model (CT26)

Achieving Exposures that Demonstrated Efficacy in all Preclinical Models Tested



Pharmacokinetic profile supports once daily dosing (14.5 hours half-life)

Doses ≥ 1.5 mg achieved exposures that demonstrated efficacy in all preclinical models tested

RXC004 Monotherapy is Safe and Well Tolerated up to 2mg



Presented at ESMO '21

Number of Patients (%)	0.5mg cohort N=4	1.0mg cohort N=3	1.5mg cohort N=7	Phase 2 dose		3.0mg cohort N=4	10.0mg cohort N=1	TOTAL N=25
				2.0mg cohort N=6				
Any TRAEs*	4 (100)	3 (100)	5 (71)	5 (83)		4 (100)	1 (100)	22 (88)
Fatigue	2 (50)	1 (33)	4 (57)	3 (50)		2 (50)	1 (100)	13 (52)
Nausea	1 (25)	2 (67)	3 (43)	1 (17)		3 (75)	1 (100)	11 (44)
Decreased appetite	1 (25)	1 (33)	2 (29)	2 (33)		3 (75)	1 (100)	10 (40)
Dysgeusia	0 (0)	0 (0)	2 (29)	4 (67)		3 (75)	1 (100)	10 (40)
Vomiting	0 (0)	1 (33)	2 (29)	1 (17)		1 (25)	1 (100)	6 (24)

- Dose limiting toxicities in 4 patients: diarrhoea, colitis and fragility fractures (10mg); colitis (3mg); ileitis (3mg); pancreatitis (2mg)
- Lower doses and denosumab prophylaxis averted bone toxicity associated with Wnt inhibition - no rises of bone turnover marker β CTX and no spontaneous fractures
- Dysgeusia appears to be dose related and did not lead to discontinuations

* All treatment related adverse events (any grade) occurring in at least 20% patients

Redx internal data

ESMO: European Society for Medical Oncology; TRAE: Treatment related adverse event; Wnt: Wingless/integrated