



Redx

Discovering Targeted Medicines

Developing Novel, Targeted Therapeutics for Cancer and Fibrotic Diseases

Corporate Presentation

November 2022

AIM:REDX



Disclaimer



THIS PRESENTATION AND ITS CONTENTS ARE NOT FOR RELEASE, PUBLICATION OR DISTRIBUTION, IN WHOLE OR IN PART, DIRECTLY OR INDIRECTLY, IN OR INTO OR FROM THE UNITED STATES OF AMERICA (THE "U.S.") (EXCEPT TO QIBS AND AIS (AS DEFINED BELOW)), CANADA, AUSTRALIA, JAPAN OR THE REPUBLIC OF SOUTH AFRICA OR ANY JURISDICTION WHERE SUCH DISTRIBUTION IS UNLAWFUL.

This presentation has been prepared by Redx Pharma plc (the "Company") solely for your information and for use at a presentation for the purpose of providing background information on the Company, its business and the industry in which it operates, and has not been approved by the United Kingdom Financial Conduct Authority, London Stock Exchange plc, the U.S. Securities and Exchange Commission or otherwise. For the purposes of this notice, "presentation" means this document, any oral presentation, any question and answer session and any written or oral material discussed or distributed during the presentation meeting.

This presentation may not be copied, distributed, reproduced or passed on, directly or indirectly, in whole or in part, or disclosed by any recipient, to any other person (whether within or outside such person's organisation or firm) or published in whole or in part, for any purpose or under any circumstances.

The presentation has not been independently verified and no representation or warranty, express or implied, is made or given by or on behalf of the Company or any of its parent or subsidiary undertakings, or the subsidiary undertakings of any such parent undertakings, or any of such person's respective directors, officers, employees, agents, affiliates or advisers, as to, and no reliance should be placed on, the accuracy, completeness or fairness of the information or opinions contained in this presentation and no responsibility or liability is assumed by any such persons for any such information or opinions or for any errors or omissions. All information presented or contained in this presentation is subject to verification, correction, completion and change without notice. In giving this presentation, none of the Company or any of its parent or subsidiary undertakings, or the subsidiary undertakings of any such parent undertakings, or any of such person's respective directors, officers, employees, agents, affiliates or advisers, undertakes any obligation to amend, correct or update this presentation or to provide the recipient with access to any further information that may arise in connection with it.

Neither this presentation nor anything contained herein shall form the basis of or be relied upon in connection with, or act as an inducement to enter into, any investment activity. This presentation does not purport to contain all of the information that may be required to evaluate any investment in the Company or any of its securities and should not be relied upon to form the basis of, or be relied on in connection with, any contract or commitment or investment decision whatsoever. This presentation is not intended to provide complete disclosure upon which an investment decision could be made. The merit and suitability of an investment in the Company should be independently evaluated and any person considering such an investment in the Company is advised to obtain independent advice as to the legal, tax, accounting, financial, credit and other related advice prior to making an investment.

To the extent available, the data contained in this presentation has come from official or third party sources. Third party industry publications, studies and surveys generally state that the data contained therein have been obtained from sources believed to be reliable, but that there is no guarantee of the accuracy or completeness of such data. While the Company believes that each of these publications, studies and surveys has been prepared by a reputable source, the Company has not independently verified the data contained therein. In addition, certain of the data contained in this presentation come from the Company's own internal research and estimates based on the knowledge and experience of the Company's management in the market in which the Company operates. While the Company believes that such research and estimates are reasonable and reliable, they, and their underlying methodology and assumptions, have not been verified by any independent source for accuracy or completeness and are subject to change without notice. Accordingly, undue reliance should not be placed on any of the data contained in this presentation.

This presentation is made available for information purposes only and does not, and is not intended to, constitute an offer to sell or an offer, inducement, invitation or commitment to purchase or subscribe for any securities, including, but not limited to, the ordinary shares of nominal value £0.01 each in the capital of the Company (the "Ordinary Shares"). The distribution of this presentation may, in certain jurisdictions, be restricted by law and neither it, nor any part of it nor the fact of its distribution shall form the basis of or be relied upon in connection with any contract and it does not constitute a recommendation regarding any securities.

Subject to certain limited exceptions, neither this presentation nor any copy of it may be taken, transmitted or distributed, directly or indirectly, into the U.S., its territories or possessions. This presentation is being provided for information purposes only and may only be made available to persons that are (i) "qualified institutional buyers" ("QIBs"), as defined in Rule 144A under the U.S. Securities Act of 1933, as amended (the "U.S. Securities Act"), (ii) "accredited investors" ("AIs") as defined in Rule 501(a) of Regulation D under the U.S. Securities Act or (iii) non-U.S. persons outside the U.S., in compliance with Regulation S under the U.S. Securities Act. By attending this presentation or by reading the presentation slides, you warrant and acknowledge that you fall within one of the categories (i), (ii) and (iii) above. Any failure to comply with the foregoing restrictions may constitute a violation of U.S. securities laws. The Company's Ordinary Shares have not been registered under the U.S. Securities Act or under any securities laws of any state of the U.S. and such securities may not be offered or sold in or into the U.S. except pursuant to the registration requirements of such act or an exemption therefrom, or in a transaction not subject to, the registration requirements of the Securities Act.

This document is limited to and intended for distribution in the United Kingdom only to persons who (i) are qualified investors (within the meaning of Article 2 of Regulation (EU) 2017/1129 as it forms part of domestic law by virtue of the European Union (Withdrawal) Act 2018) and (ii) who have professional experience in matters relating to investments who fall within the definition of "investment professionals" in Article 19(5) of the Financial Services and Markets Act 2000 (Financial Promotion) Order 2005 (the "FPO") and/or to high net worth bodies corporate, unincorporated associations and partnerships and trustees of high value trusts as described in Article 49(2)(a) to (d) of the FPO or to those persons to whom it may otherwise be lawfully communicated (in each case referred to as "Relevant Persons"). This document has not been approved by an authorised person for the purposes of Section 21 of the Financial Services and Markets Act 2000. The information contained in this document is not intended to be viewed by, or distributed or passed on (directly or indirectly) to, and should not be acted upon by any class of persons other than Relevant Persons. Any investment or investment activity to which this document relates is available only to Relevant Persons and will be engaged only with Relevant Persons.

This document is limited to and intended for distribution in the European Economic Area only to persons who are qualified investors (within the meaning of Article 2 of Regulation (EU) 2017/1129).

This presentation includes forward looking statements. The words "expect", "anticipate", "intends", "plan", "estimate", "aim", "forecast", "project" and similar expressions (or their negative) identify certain of these forward looking statements. These forward looking statements are statements regarding the Company's intentions, beliefs or current expectations concerning, among other things, the Company's results of operations, financial condition, liquidity, prospects, growth, strategies and the industry in which the Company operates, and include statements regarding the Company's ongoing and planned clinical trials, regulatory approval process, and demand for the Company's product candidates which are subject to risks, uncertainties, and other factors that could cause actual results to differ materially from those suggested by forward-looking statements. These factors include, but are not limited to, the following: the Company has incurred losses since its inception and anticipates that it may continue to incur losses for the foreseeable future; the Company does not expect to generate any material income until its pipeline of programmes are progressed commercially; the Company will need to raise additional funding in the future, which may not be available on acceptable terms, or at all; and the Company may not be able to obtain exclusivity or intellectual property rights for its product candidates or prevent others from developing similar competitive products. The forward looking statements in this presentation are based on numerous assumptions regarding the Company's present and future business strategies and the environment in which the Company will operate in the future. Forward looking statements involve inherent known and unknown risks, uncertainties and contingencies because they relate to events and depend on circumstances that may or may not occur in the future and may cause the actual results, performance or achievements of the Company to be materially different from those expressed or implied by such forward-looking statements. Many of these risks and uncertainties relate to factors that are beyond the Company's ability to control or estimate precisely, such as future market conditions, currency fluctuations, the behaviour of other market participants, the actions of regulators and other factors such as the Company's ability to continue to obtain financing to meet its liquidity needs, changes in the political, social and regulatory framework in which the Company operates or in economic or technological trends or conditions. Past performance should not be taken as an indication or guarantee of future results, and no representation or warranty, express or implied, is made regarding future performance. The Company expressly disclaims any obligation or undertaking to release any updates or revisions to these forward looking statements to reflect any change in the Company's expectations with regard thereto or any change in events, conditions or circumstances on which any statement is based after the date of this presentation or to update or to keep current any other information contained in this presentation. Accordingly, undue reliance should not be placed on the forward looking statements, which speak only as of the date of this presentation.

By attending the meeting where this presentation is made or by accepting a copy of this presentation, you agree to be bound by the foregoing limitations and to maintain absolute confidentiality regarding the information disclosed in this presentation. Without limiting the generality of the foregoing, the recipient acknowledges and agrees that (i) photocopying or other duplication or transmittal of the presentation in whole or in part and the data provided without prior written consent of the Company is strictly prohibited; and (ii) the Company will not be subject to any liability based on the information contained in this presentation, errors herein or omissions here from, whether or not the Company knew or should have known of any such errors or omissions, and/or was responsible for and/or participated in such error or omission from this presentation.

Our ambition: To create world leading medicines to transform patients' lives



Lisa Anson
CEO

High profile general manager, former President of AstraZeneca UK, with >25 years in biotech and pharma



Dr Richard Armer
CSO

Accomplished drug discovery executive, with >25 years in biotech and pharma



Dr Jane Robertson
CMO

Oncologist with >18 years in biotech and pharma drug development. Led the successful clinical development of Lynparza



Peter Collum
CFO

Experienced finance and strategy executive with >25 years in biopharma



Developing Novel Targeted Medicines for High Unmet Medical Needs



Developing novel, small molecule, highly targeted medicines for cancer and fibrotic disease, and the emerging area of cancer-associated fibrosis



Clinical-stage biotech with two Phase 2 assets against compelling clinical targets, and Discovery Engine fuelling pipeline

Porcupine

ROCK

Experienced scientific team leveraging distinct approach. Track record of consistently generating differentiated drug candidates

5 clinical molecules discovered


Backed by blue chip specialist biotech investors. Cash runway into 2024 to deliver multiple value-inflection points

Redmile Group

SOFINNOVA
PARTNERS



Validated Discovery Engine Fuels Pipeline



ADME: Absorption, distribution, metabolism and excretion; IND: Investigational new drug application; PoC: Proof of concept; PK: Pharmacokinetic

Advancing a Robust Pipeline Built In-House



	Target/ Product	Indication(s)	Research	Preclinical	Phase 1	Phase 2	Phase 3	Upcoming Milestones
Oncology	Porcupine Inhibitor (RXC004)	Genetically selected MSS mCRC monotherapy + anti-PD-1 combination	PORCUPINE					Topline data - from H1 2023
		Genetically selected pancreatic cancer monotherapy Biliary cancer monotherapy + anti-PD-1 combination	PORCUPINE2					
Fibrosis	ROCK2 Selective Inhibitor (RXC007)	Idiopathic pulmonary fibrosis (IPF)						Phase 2a topline data - H2 2023
	GI-targeted ROCK Inhibitor (RXC008)	Fibrostenotic Crohn's disease						IND/CTA submission - end 2023
Discovery Engine	DDR Inhibitor (Discoidin Domain Receptor)	Fibrosis, cancer-associated fibrosis						Progress programmes - target of 2 INDs by 2025
	Research Targets (Multiple Programmes)	Oncology & fibrosis						
Partnered	Porcupine Inhibitor (RXC006/AZD5055)	Idiopathic pulmonary fibrosis (IPF)						Licensed to AstraZeneca
	Pan-RAF Inhibitor (JZP815)	Oncology						Sold to Jazz Pharmaceuticals
	MAPK Pathway Target	Oncology						Progress Jazz collaboration

RXC007: A Selective ROCK2 Inhibitor for Fibrotic Diseases - Lead Indication is IPF with Phase 2 Data Expected H2 2023




Highlights

- RXC007 is a highly potent, selective and orally-active ROCK2 inhibitor
- ROCK2 is a validated, compelling target at a key junction in cell signalling pathways central to fibrosis
- Robust preclinical efficacy data across disease models supports clinical development plan in lung fibrosis – IPF and CF-ILD
- Phase 1 data in SAD and multi-dose cohorts confirms drug like profile for safety and PK
 - No significant adverse events observed in healthy volunteers and predictable PK profile
- **Phase 2a in IPF, expected to report H2 2023**, is a dose-ranging study to inform Phase 2b dose
 - Initial 12-week Phase 2a study for early efficacy readouts, safety and tolerability in IPF patients +/- SoC, in addition to target and disease biomarker engagement
- Phase 2b planned for RXC007 plus SoC over 12 months with lung function (FVC) as primary endpoint

CF-ILD: Chronic fibrosing interstitial lung disease; FVC: Forced vital capacity; IPF: Idiopathic pulmonary fibrosis; PK: Pharmacokinetic; ROCK: Rho associated protein kinase; SAD: Single ascending dose; SoC: Standard of care

ROCK is a Compelling, Nodal Target for Fibrotic Diseases



Why ROCK

- 1** RhoA/ROCK/ROCK2 downstream of many major profibrotic factors
- 2** ROCK is involved in diverse cellular processes
- 3** ROCK upregulates key profibrotic genes. Upregulation of these genes leads to actin cytoskeleton organisation, cell adhesion and motility, proliferation, and extra cellular matrix remodeling

Why ROCK2

- The role of ROCK2 in a diverse range of cellular process allows RXC007 to have pleiotropic effects
- Systemic inhibition of ROCK1&2 results in hypotension, not seen with selective ROCK2 inhibition
- ROCK2 inhibition alone is sufficient to protect from pulmonary fibrosis in mouse models⁽¹⁾

(1) Knipe, 2018


Highly Selective ROCK2 Inhibition Avoids Hypotension and Limits Off-target Toxicity



- Selectively inhibiting ROCK2 has been notoriously challenging
- High target selectivity desirable to limit off target toxicity
- RXC007 is potent and highly selective
 - >100-fold versus ROCK1 in biochemical assays
 - >100-fold selectivity against 468 kinases
- RXC007 Phase 1 in healthy volunteers showed
 - No SAEs
 - All AEs transient, mild and reversible with no dose changes required
 - No evidence of hypotension

RXC007 is highly selective against ROCK1 and other kinases

Plot of kinases inhibited by RXC007 with $IC_{50} < 1 \mu M$



- IC_{50} determination at K_m ATP of the 20 kinases inhibited with >50% following 468-kinase panel at $10 \mu M$
- Only 2 kinases inhibited at $IC_{50} < 1 \mu M$
 - ROCK2 the most potently inhibited kinase ($0.004 \mu M$)
 - MEK2 only off-target below $1 \mu M$ ($0.428 \mu M$)

Lung Fibrosis: A Growing Clinical Need with Limited Treatment Options




Lead indication is Idiopathic Pulmonary Fibrosis (IPF)


- IPF is a deadly fibrotic lung disease
- Median survival between 3 – 5 years⁽¹⁾
- High unmet need: over 170,000+ patients in the US, EU5 and Japan alone
- Global IPF market projected to reach \$3.6 billion by 2029⁽²⁾
- Nintedanib and pirfenidone are the only approved treatments but have significant side effects that limit use

Broader opportunity in Interstitial Lung Disease (ILD)

- IPF represents approx. one third of ILD patients⁽³⁾
- Of the two thirds of ILD patients without IPF, up to 40% may develop a progressive fibrosing phenotype⁽⁴⁾ (PF-ILD) where new anti-fibrotic treatments are needed
- PF-ILD includes rheumatoid arthritis-ILD (RA-ILD), systemic sclerosis-ILD (SSc-ILD) and mixed connective tissue disease-ILD (mCTD-ILD)
- SSc-ILD market is particularly well-characterized, with this segment alone estimated to surpass \$1.1 billion by 2030⁽⁵⁾



Normal lung



Fibrotic lung

(1); (2) Patient numbers (diagnosed prevalence) & market size forecast data sourced from Global Data (US, EU5, Japan)


(3) Kaul 202 Clinical Estimates from Hyun 2015, Ley 2012, Raghu 2006 (4) Olson 2021; (5) Clarivate, Key Findings on Scleroderma (Systemic Sclerosis) Published: September 2021

Strong Literature Rationale on ROCK and ROCK2 Supports IPF as a Lead Indication



ROCK as a target in IPF⁽¹⁾


RhoA/ROCK/ROCK2 signalling pathway is upregulated in human IPF patients



(Myo)fibroblasts isolated from lungs of patients with IPF (n = 8), non-IPF control subjects (n = 6) ROCK activity assays using a colorimetric approach. Data are mean ± SD

ROCK2 expression in IPF⁽²⁾


Increased ROCK2 expression in IPF bronchial epithelial cells and fibroblasts



Intensity of immunostaining evaluated on a scale of 0 to 3
N=6 per group; NF: not found

Preclinical validation of selective ROCK2 inhibition⁽³⁾

ROCK2+/- knockout mice are protected from bleomycin-induced pulmonary fibrosis




Genetic reduction in either or both ROCK isoforms affords protection from bleomycin-induced pulmonary fibrosis
Significant reduction in hydroxyproline content (lung collagen)

(1) Zhou, 2013; (2) Shimizu, 2014; (3) Knipe, 2018

Preclinical Efficacy Data Demonstrates Pleiotropic Effects of Selective ROCK2 Inhibition Supporting IPF as Lead Indication


Therapeutic Murine Bleomycin-induced Lung Fibrosis Model (BLM)

Reduction of collagen deposition




Reduction of fibrosis

Representative images
Lung histology Sirius red




Vehicle



RXC007 50 mg/kg BID

Suppression of the expression of genes strongly associated with fibrosis in IPF




Mouse 21-day therapeutic model (dosing from day 7)
Data generated by Redx

RXC007 Modulates Pro-Fibrotic Gene and Protein Expression in *Ex-vivo* Lung Tissue from Patients with Pulmonary Fibrosis



Methodology : Analysis of RXC007 in live IPF tissue samples

Cores of tissue from 3 resected ILD lungs and cultured *in vitro*




Compound addition to supernatant for 48 hours

- RXC007: 0.3, 3, 30µM
- Nintedanib: 0.1, 1µM


Tissues analysed for pro-fibrotic gene expression
Supernatants analysed for pro-fibrotic proteins
Tissue proven to be live
Compounds shown to be non-toxic

RXC007 modulated inflammation & tissue remodelling gene expression pathways, similar profile to nintedanib

Number of pathways significantly increased or decreased out of 103 measured



RXC007 suppressed expression of pro-fibrotic proteins (examples)




Notes: Gene expression analysis was analysed using KEGG pathway grouping and Z-scores to normalise analysis across donors. RXC007 was shown to suppress expression of 20 out of 24 protein analytes analysed from the supernatant at 30µM but not at 3 or 0.3µM. Nintedanib did not suppress protein analytes at either concentration
Data generated by Redx

ILD: Interstitial lung disease


Proof-of-Concept Demonstrated in an Immune Mediated Model

Murine Sclerodermatous chronic Graft versus Host Model

Skin thickness




Hydroxyproline skin




Syngeneic
 (A) + RXC007 5 mg/kg BID
 (A) + Nintedanib 30 mg/kg QD
 Allogeneic (A)
 (A) + RXC007 50 mg/kg BID
 (A) + Nintedanib 50 mg/kg QD

Data generated by Redx, as presented at ICLAF 2022





Cutaneous cGVHD Score



Significant Reduction of Fibrosis Score and Collagen Content in the Lungs Supports Broader Development in Lung Fibrosis



Murine Sclerodermatous chronic Graft versus Host Model




□ Syngeneic
■ Allogeneic
□ Allogeneic + RXC007 5 mg/kg BID
■ Allogeneic + RXC007 50 mg/kg BID
□ Allogeneic + Nintedanib 30 mg/kg QD
■ Allogeneic + Nintedanib 50 mg/kg QD

Data generated by Redx, as presented at ICLAF 2022

Phase 1 Data in Healthy Volunteers Showed Good Safety and Pharmacokinetic Profile



Multidose cohort 50 mg BID - total plasma concentration over time



Good safety profile

- No SAEs reported in SAD and multidose cohorts
- Safe and well tolerated with few treatment emergent adverse events reported
- All AEs transient, mild and reversible with no dose changes required
- No evidence of hypotension, validating rationale for selective ROCK2 inhibition

Drug-like pharmacokinetic profile


- Pharmacokinetics as predicted from preclinical data
- Mean half-life 9-11 hours, potentially suitable for once-daily dosing
- Essentially linear exposure in SAD from 2 mg QD to 70 mg BID
- No significant differences between 50 mg fed and fasted cohorts
- Achieved biologically relevant exposures from 20 mg BID and above based on preclinical models

PK sampling up to 72 h; only 0-24 h plotted
Data generated by Redx, as presented at ICLAF 2022

BID: Twice daily; QD: Once daily; SAD: Single ascending dose; SAE: Serious adverse event

Phase 2a dose ranging study to inform Phase 2b dose

Provides early efficacy readouts, safety and tolerability in IPF patients with or without standard IPF therapy



DLCO: Carbon monoxide diffusion coefficient; FVC: Forced vital capacity; HRCT: High resolution computerised tomography; Pbo: Placebo; PK: Pharmacokinetics

RXC004: Clinically Validated Porcupine Inhibitor in Phase 2




Highlights

- RXC004 is a highly potent, orally active, once daily Porcupine inhibitor
- Porcupine inhibition blocks secretion of all Wnt ligands, preventing both tumour growth and immune evasion
- RXC004 demonstrated clinical target engagement at all doses and has optimal PK profile with once daily, oral dosing
- RXC004 was well tolerated in Phase 1, both monotherapy and in combination with nivolumab, and achieved exposures clinically that demonstrated efficacy in pre-clinical models
- RXC004 demonstrated differential clinical efficacy in Wnt-ligand dependent tumours in Phase 1 monotherapy cohort (ESMO 2021)
- Combination with anti-PD-1 treatment could open new patient segments e.g. MSS mCRC, and anti-PD-1 resistant patients
- Recruitment ongoing, Phase 2 programme in Wnt-ligand dependent tumours – **data expected from H1 2023**
 - In genetically selected Wnt-ligand driven tumours, monotherapy and anti-PD-1 combination settings
 - In tumours with high Wnt-ligand expression, monotherapy and anti-PD-1 combination settings
 - Phase 2 dose selected for monotherapy 2 mg, and 1.5mg in combination with anti-PD-1
 - Media reported single biliary cancer patient partial response at six months

Porcupine Inhibition Targets Wnt-Ligand Dependent Tumours



Wnt signalling pathway is well established as a critical driver of cancer




- 1** Porcupine is a key enzyme upstream in the Wnt pathway, which adds a lipid chain to all 19 Wnt ligands (palmitoylation) enabling secretion
- 2** Wnt ligands activate both canonical and non-canonical signalling pathways
- 3** Canonical and non-canonical target genes drive tumour growth and tumour immune evasion
- 4** By inhibiting Porcupine, RXC004 blocks the release of all Wnt ligands from cells **inhibiting tumour growth** and **reversing immune evasion**
- 5** Targeting Wnt-ligand dependent tumours provides a clear **patient selection strategy**. Tumours with RSPO fusion or RNF43 mutations see upregulation of the Wnt pathway (increased number of frizzled receptors, through impaired degradation). Beyond genetic mutations, some other tumours display elevated Wnt-ligand expression

RNF43: Ring finger protein 43; RSPO: R-spondin; Wnt: Wingless/integrated


Patient Selection Strategy: Wnt-Ligands can Drive Growth and Immune Evasion in the Tumour Microenvironment




Genetic aberrations lead to Wnt-ligand driven tumour growth



Wnt-ligand expression can drive tumour growth




Wnt-ligands negatively impact immune cell function, resulting in immune evasion and cold tumours



MDSC: Myeloid-derived suppressor cells; RNF43: Ring finger protein 43; RSPO: R-spondin; Wnt: Wingless/integrated

Dual Mechanism of Action Demonstrated Preclinically


Inhibits tumour cell proliferation



RXC004 inhibits tumour growth in an RNF43 mutant human pancreatic cancer model


DIRECTLY TARGETS TUMOUR CELL GROWTH

Alters tumour cell metabolism



RXC004 treated tumour cells are less metabolically active. Cells also change morphology and increase mucous secretion


As a monotherapy



Increased survival rate post-RXC004 monotherapy treatment in an anti-PD-1 resistant 'cold' mouse melanoma syngeneic model (B16F10)

REVERSES WNT PATHWAY INDUCED IMMUNE EVASION

In combination with anti-PD-1



Combination of RXC004 + anti-PD-1 showed a statistically relevant change in ratio of Cytotoxic T cells: Regulatory T cells, compared to either monotherapy alone in PD(L)-1 axis dominated CT26 CRC model

Data as of 7th June 2021
Source: Redx posters, <https://aacrjournals.org/cancerrescommun/article/2/9/914/708958/The-Wnt-Pathway-Inhibitor-RXC004-Blocks-Tumor>

CRC: Colorectal cancer; RSPO: R-Spondin

High Unmet Need in Hard-to-Treat Wnt-Ligand Dependent Tumours



Wnt-ligand dependence mechanism	Tumour	5-year survival rate ⁽¹⁾	Prevalence of Wnt-ligand dependence	Annual Wnt-ligand dependence incidence ⁽⁶⁾	Approach being tested in Ph2	
					Monotherapy	anti-PD-1 Combo
Wnt-ligand driven	MSS mCRC 	14%	8% RNF43 mutation & RSPO fusion ⁽²⁾⁽³⁾	14,000 patients	✓	✓
	Pancreatic 	3%	7% RNF43 mutation ⁽⁴⁾	9,000 patients	✓	✗
Wnt-ligand expressed	Biliary 	2%	70% Wnt-ligand expression ⁽⁵⁾	51,000 patients	✓	✓

LCM potential in multiple additional tumours

Monotherapy - opportunity to pursue a tumor agnostic approach - RNF43 mutations and RSPO fusions also detected in MSI-H mCRC, CRPC and squamous NSCLC

Combination - opportunity to target any anti-PD-1 resistant tumour with high Wnt-ligand expression (~28 tumour types)

(1) www.cancer.net (2) RNF43 mutation frequency determined from all relevant studies published on cBioPortal for cancer genomics (updated Jan 2018). Only mutations resulting in functional impairment (LoF) were considered. Gao et al. 2013 & Cerami et al. 2012. (3) RSPO fusion prevalence in CRC is a combination of studies (Shesagiri, 2012; Shinmura, 2014; Kleeman, 2019); Redx notes the combined figure is an approximate figure calculated on the basis of RNF43 mutations and RSPO fusions based on the studies; indicated and these studies included a limited dataset of samples; (4) Witkiewicz 2015; (5) Loilome et al. 2014, Boulter et al. 2015; (6) Incidence data sourced from GlobalData Epidemiology data (Major Markets: US, EU5, Japan, China)

RXC004 was Safe and Well Tolerated in Monotherapy and in Combination with anti PD-1 at Phase 2 Doses



Phase 1 Study Data	1.0mg monotherapy N=3	1.0mg nivo combination N=5	1.5mg monotherapy N=7	Phase 2 doses		3.0mg monotherapy N=4	TOTAL N=33*
				1.5mg nivo combination N=8	2.0mg monotherapy N=6		
Any TRAEs*	3 (100)	5 (100)	5 (71)	8 (100)	5 (83)	4 (100)	30 (91)
Fatigue	1 (33)	2 (40)	4 (57)	3 (38)	3 (50)	2 (50)	15 (45)
Nausea	2 (67)	2 (40)	3 (43)	4 (50)	1 (17)	3 (75)	15 (45)
Dysgeusia	0 (0)	1 (20)	2 (29)	4 (50)	4 (67)	3 (75)	14 (41)
Decreased appetite	1 (33)	1 (20)	2 (29)	3 (38)	2 (33)	3 (75)	12 (36)


- The recommended Phase 2 dose of RXC004 is 2mg in monotherapy
- Both RXC004 1.0mg and 1.5mg doses were safe and tolerable in combination with nivolumab
 - RXC004 doses higher than 1.5mg were not explored in Module 2 because of the potential for overlapping toxicity of colitis, which was reported in RXC004 monotherapy Module 1 and is a known adverse effect of immune checkpoint inhibitors.

TRAE: Treatment related adverse event

Phase 1 Clinical Efficacy Data Supports Development in Wnt-Ligand Dependent Tumours



Phase 1 Monotherapy All Comers Study Clinical Activity by Wnt-Ligand Dependence†



Numbers= dose in mg
 † Study was in unselected patients; retrospective analysis
 * RNF43 LoF mutation # RSPO Fusion
 Data generated by Redx: Monotherapy Data cut-off date 30 July 2021, as presented at ESMO 2021; Combination Data Cut off 18 Oct 2022, as presented at SITC 2022

Phase 1 monotherapy efficacy data

18/25 monotherapy patients had RECIST-evaluable disease

5/7 patients with Wnt-ligand dependent tumours achieved durable RECIST Stable Disease (SD)

- In addition, 2/7 patients with Wnt-ligand dependent tumours had SD in target lesions but disease progression overall
- 0/11 patients with unknown or Wnt-ligand independent tumours had SD

Median treatment duration higher in patients with Wnt-ligand dependent tumours

- 13.1 weeks in Wnt-ligand dependent tumours
- 6.6 weeks in unknown or Wnt-ligand independent tumours

Phase 1 combination efficacy data

10/13 patients had RECIST evaluable disease

- 4/6 patients with unselected tumours in the 1.5mg cohort had RECIST stable disease as best response

APC: Adenomatous polyposis coli gene; CRC: Colorectal cancer; LoF: Loss of function; PD: Progressive disease; RECIST: Response evaluation criteria in solid tumours; RNF43: Ring finger protein 43; RSPO: R-spondin; SD: Stable disease; Wnt: Wingless/integrated

Phase 2 Programme in Wnt-Ligand Dependent Tumours Expected to Deliver Topline Data During 2023



	PORCUPINE MSS Metastatic Colorectal Cancer		PORCUPINE2 Pancreatic and Biliary Cancer		
Endpoints	Monotherapy Genetically Selected (n=20)	Combination Genetically Selected RXC004 + anti-PD-1 (n=20)	Monotherapy Genetically Selected Pancreatic Cancer (n=15)	Monotherapy Biliary Cancer (n=15)	Combination Biliary Cancer RXC004 + anti-PD-1 (n=15)
	DCR Secondary Tumour size change, ORR, DoR, PFS	ORR Secondary DCR, Tumour size change, DoR, PFS	PFS at 6 months Secondary - DCR, OS, ORR, DoR and % change in sum of target lesions		ORR Secondary DCR, Tumour size change, DoR, PFS
Topline data expected	H2 2023	H2 2023	H2 2023	H1 2023	H2 2023
Analysis	Key translational endpoints will characterise mechanism of action • FDG-PET scans to assess metabolic effect of RXC004 on tumours • Tracking circulating tumour DNA (ctDNA) • Baseline and on-treatment biopsies to assess changes to tumour and immune microenvironment				

DCR: Disease control rate; DoR: Duration of response; ORR: Overall response rate; OS: Overall survival; PFS: Progression free survival

RXC008: GI-targeted ROCK Inhibitor for Fibrostenotic Crohn's Disease Planned to Enter Phase 1 Clinical Trial End 2023



Highlights


- RXC008 is a potent, oral, small molecule non-systemic ROCK 1/2 inhibitor
- Fibrostenotic Crohn's disease is a significant unmet need - only treatment option for patients is successive surgical intervention
- RXC008 is a potential first-in class treatment - no approved therapies for underlying fibrosis and no curative treatments available
- ROCK is a key target involved in fibroblast activation, and is upregulated in fibrostenosis in Crohn's disease
- RXC008 is GI-targeted, selectively active in gut without risking systemic exposure
- RXC008 has demonstrated robust preclinical efficacy *in vivo*
- IND enabling work underway - **Phase 1 start planned for end 2023**

GI: Gastrointestinal; IND: Investigational new drug application; ROCK: Rho associated protein kinase;

Potential First-in-Class Treatment for Fibrostenotic Crohn's Disease




Clinical progression in Crohn's




1.7 million⁽¹⁾ patients globally affected by Crohn's disease

Fibrotic stricture formation



>50% of patients⁽²⁾ develop fibrostenosis and strictures within 10 years of first diagnosis


Surgical interventions



No approved therapies for underlying fibrosis only treatment options are debilitating surgical intervention

(1) Clarivate, Crohn's disease disease landscape & forecast pg 39, Published Sep 2022; (2) Chan et al, 2018

RXC008: GI-targeted pan-ROCK Inhibitor Targets a Fibrotic Pathway Nodal Point without Systemic Breakthrough




- ROCK is a nodal point in the fibrotic signalling pathway
- Inhibiting ROCK 1&2 systemically is known to result in hypotension
- GTR inhibitors are specifically designed to avoid hypotensive effects associated with systemic ROCK inhibition
- RXC008 designed to be retained in the GI tract via high efflux and low permeability, rapidly metabolised by paraoxonase enzymes in the plasma
- Result of this is virtually no systemic breakthrough

GI: Gastrointestinal; GTR: Gastrointestinal targeted ROCK; MRTF: myocardin-related transcription factor; MLC: myosin light chain


RXC008 Reduces Fibrosis in the Adoptive T-cell Transfer Model

45-day Adoptive T-cell transfer model of Crohn's disease (with therapeutic oral compound administration from day 14)

RXC008 reduces tissue damage




RXC008 reduces fibrosis





p40 data has been modified for the purpose of graphical representation
Colon damage could not be detected in the p40 treated animals.

Chronic DSS GI fibrosis model* (RXC008 dosed orally, prophylactically for 9 weeks)

Reduction in fibrosis measured by histology



Reduction in fibrosis measured by MRI





*Research in collaboration with Ghent University
1-way Anova with Dunnet's multiple comparison, # Vehicle day 0 v day 61 * vehicle day 61 v RXC008 day 61
Data generated by Redx

DSS: Dextran sulfate sodium; MT: Magnetization transfer; QD: Once daily

ROCK Inhibition Potently Suppressed or Reversed Fibrosis in Preclinical GI Models




GITR inhibition was able to reverse fibrosis in a therapeutic 12-week DSS model
 (Compound administered orally once a day from week 6 when fibrosis established)



Data generated by Redx, REDX8087 is similar to RXC008

DSS: Dextran sulfate sodium

Redx is Funded into 2024 to Deliver Significant Pipeline Progression



AIM (UK) listed Ticker: REDX
Total shares in issue: 334,911,458
Fully diluted: 482,210,019**

* At completion of funding on 8 June 2022

** Assuming full conversion of loan notes and exercise of employee share options. Updated 7 November 2022

Multiple Potential Milestones During 2022 and 2023



	2022 (CY)	2023 (CY)
Porcupine Inhibitor (RXC004)	✓ Data from Phase 1 combination study with nivolumab (anti-PD-1) ✓ Initiation of PORCUPINE2 monotherapy Phase 2 study in genetically selected pancreatic and biliary cancer ✓ Initiation of anti-PD-1 combination arms	Ph2 data - from H1 2023 PORCUPINE • H2 mono & combination - MSS mCRC PORCUPINE2 • H1 monotherapy - biliary • H2 combination - biliary • H2 monotherapy - pancreatic
ROCK2 Selective Inhibitor (RXC007)	✓ H1 Data from Phase 1 healthy volunteer study ✓ Initiation of Phase 2a study in IPF • H2 US IND	H2 Phase 2a topline data read out
GI-targeted ROCK Inhibitor (RXC008)	✓ H1 Select development candidate	IND/CTA submission – end 2023
DDR Inhibitor (Discoidin Domain Receptor)	• Progress DDR inhibitor for fibrosis • Advance multiple programmes in early discovery	Progress target of two additional wholly-owned INDs by 2025
Research Targets		
Porcupine Inhibitor (RXC006/AZD5055)		
Pan-RAF Inhibitor (JZP815)	✓ IND cleared triggering \$5m June 2022	Redx remains eligible for regulatory and development milestones
MAPK Pathway Target		


CY: Calendar year; CTA: Clinical trial application



Redx

Discovering Targeted Medicines

Appendix



Monotherapy is Safe and Well Tolerated at Phase 2 Recommended Dose




Number of Patients (%)	Phase 2 dose						TOTAL N=25
	0.5mg cohort N=4	1.0mg cohort N=3	1.5mg cohort N=7	2.0mg cohort N=6	3.0mg cohort N=4	10.0mg cohort N=1	
Any TRAEs*	4 (100)	3 (100)	5 (71)	5 (83)	4 (100)	1 (100)	22 (88)
Fatigue	2 (50)	1 (33)	4 (57)	3 (50)	2 (50)	1 (100)	13 (52)
Nausea	1 (25)	2 (67)	3 (43)	1 (17)	3 (75)	1 (100)	11 (44)
Decreased appetite	1 (25)	1 (33)	2 (29)	2 (33)	3 (75)	1 (100)	10 (40)
Dysgeusia	0 (0)	0 (0)	2 (29)	4 (67)	3 (75)	1 (100)	10 (40)
Vomiting	0 (0)	1 (33)	2 (29)	1 (17)	1 (25)	1 (100)	6 (24)

- Dose limiting toxicities in 4 patients: diarrhoea, colitis and fragility fractures (10 mg); colitis (3 mg); ileitis (3 mg); pancreatitis (2 mg)
- Lower doses and denosumab prophylaxis averted bone toxicity associated with Wnt inhibition – no rises of bone turnover marker β CTX and no spontaneous fractures
- Dysgeusia appears to be dose related and did not lead to discontinuations

Redx internal data
 * All treatment related adverse events (any grade) occurring in at least 20% patients

ESMO: European Society for Medical Oncology; TRAE: Treatment related adverse event; Wnt: Wingless/integrated


MSS mCRC Patients Have Poor Outcomes with Current Standards of Care



(1) Aparicio et al 2020 (2) Tabernero 2022 (3) (4) Kopetz 2021 (Beacon) Giordano et al, 2021(5) Vegal et al, 2016

DCR: Disease control rate; MSI: Microsatellite-unstable
ORR: Overall response rate; OS: Overall survival; PFS: Progression free survival; RNF43: Ring finger protein 43; RSPO: R-spondin; Wnt: Wingless/integrated


Currently, 2% Overall Survival Rate for Biliary Tract Cancers, Opportunity in Late Line Settings to Reverse Immune Evasion



*N.B. FDA approval of durvalumab + chemo received by AstraZeneca in September 2022. Pembrolizumab Phase 3 ongoing with commercial use subject to regulatory approval.
(1) Valle et al NEJM 2010; (2) Oh et al NEJM Evidence 2022; (3) Lamarca et al Lancet 2019

BTC: Biliary tract cancer; Wnt: Wingless/integrated

Discoidin Domain Receptor (DDR) Inhibition as a Potential Novel Therapeutic Class for Fibrosis



DDR is a collagen target

- Non-integrin tyrosine kinase collagen receptors
- Collagen binding initiates downstream fibrotic signalling pathways

DDR inhibition is a novel approach

- Novel, druggable therapeutic target for fibrosis
- Recent publication of preclinical PoC studies for small molecule inhibitors in models of lung and kidney fibrosis
- Potential for disease modifying efficacy

Redx has a discovery programme

- Potent and selective DDR inhibitors identified
- In lead optimisation phase
- Preclinical PoC ongoing in lung and kidney fibrosis models